GENERAL INDEX TO VOLUME XXXV

New scientific names of plants and the final members of new combinations are printed in bold-face type; synonyms and page numbers having reference to figures and plates, in italics; and all other matter in ordinary type.

A

Absorption spectra of epidermal cells from a Coleus leaf, 243

Acosta, Joseph de, description of maize by, 151, 153, 154

Adenocalymma bracteatum, calyx glands of, 132

Allen, Caroline K. Lauraceae of Panama, 1 American carboniferous floras, Contributions to our knowledge of, 193

Amsonia, a new, from the Trans-Pecos,

Amsonia hirtella, 238; Palmeri, 238; Peeblesii, 238; Tharpii, 237

Anacardiaceae, living, compared with a fossil wood, 109, 110

Anderson, Edgar: and William L. Brown. A morphological analysis of row number in maize, 323; William L. Brown and. The southern dent corns, 255

Andrews, Henry N.: A note on Fomes idahoensis Brown, 207; and Robert W. Baxter. Contributions to our knowledge of American carboniferous floras. X. An Osmundaceous stem from Iowa, 193; and Sergius Mamay. A Crossotheca from northern Illinois, 203

Anemopaegma Chamberlaynii, glandular fields in leaflet of, 130, 136; glandular scale on, 134

Anthocyanin concentration in hair cells of Coleus leaf, 244, 249, 252

Apocynaceae and Asclepiadaceae, Miscellaneous new, 233

Arrabidaea Blanchetii, interpetiolar glands of, 128; obliqua, domatia on leaflet, 130 Asclepiadaceae, Miscellaneous new, 233

B

Bauhin, Caspar and Jean, herbals of, 165, 178; illustration of maize in, 178

Baxter, Robert W.: A study of the vegetative anatomy of the genus Sphenophyllum from American coal balls, 209; Henry N. Andrews and. Contributions to our knowledge of American Carboniferous floras. X. An Osmundaceous stem from Iowa, 193

Beilschmiedia, 109; roxburghiana, 110 Bignoniaceae, The uses of glands in a taxonomic consideration of the family, 123 Bischofia javanica, 110

Boccone, herbal of, 178; illustration of maize in, 178

Bock, Hieronymus, account of maize in herbal of, 152, 156, 159

Bonnett, O. T. Ear and tassel development in maize, 269

Bridelia minutiflora, 109, 110

Brown, William L.: and Edgar Anderson. The southern dent corns, 255; Edgar Anderson and. A morphological analysis of row number in maize, 323

Brunfels' herbal, 159

Burseraceae, living, compared to a fossil wood, 109, 110

C

Callichlamys latifolia, calyx glands on leaflet of, 136

Camerarius, account of maize in herbal of, 162, 176; illustration in, 166

Canarium, 109; rufum, 110

Capparidaceae, 139; of Panama, 75

Carboniferous floras, Contributions to our knowledge of. X. An Osmundaceous stem from Iowa, 193

Castilla, 119

Cells, physiological activity of, 239

Ceratophytum tobagense, neuro-axillary glands of, 130

Cleome, 139; brachycarpa, 143, 144; decipiens, 139, 142; densiflora, 144, β pallens, 144; gracilis, 144, β turgescens, 144; birsuta, 144; lateralis, 144; Macrothyrsus, 144; porphyrantha, 144; puberula, 144

Coal balls, American: fossil plants in, 209; from Iowa, 193

Coleus: pigment production in hair cells, 243, 243; absorption spectra for cells from a red and green, 244, 245-248, 251

Columbus's report on maize, 156
Commoner, Barry. Quantitative determination of the pigment content of single
cells by means of a new microspectrophotometer, 239

Conference, the 1947, on the inflorescences of Zea Mays, 269

Contributions to our knowledge of American carboniferous floras. X. An Osmundaceous stem from Iowa, 193

(407)

Cordus, description of maize in herbal of,

Corn Conference of November, 1947, papers read at, 269-406

Corn: breeding, 267; cob, anatomy of, 353; embryos, Right-handed and left-handed, 317; four-rowed ear, Vascular anatomy of, 337, 340; Indian and Turkish, 173-176; southern dent varieties, 267

Corn, see also Maize and Zea Mays Corn grass, 309, 310

Corns, The southern dent, 255

Crossotheca from northern Illinois, A, 203 Crossotheca, 203; Boulayi, 204; communis, 204; Hoeninghausi, 204; McLuckiei, 203, 204; pinnatifida, 204; sagittata, 204; Schartzlarensis, 204; trisecta, 204

Cruciferae of Panama, 99

Cunuria, 118

130

Cutler, Hugh C., and Marian C. Studies in the structure of the maize plant, 301 Cydista heterophylla, glandular fields in,

Dent corns, the southern, 255

Dicotyledonous wood found associated with the Idaho Tempskyas, 107

Dodonaeus, description of maize in herbal of, 162, 173; illustrations of maize in, 163

Domestication: of Hevea, 117; of maize, 394

Dothidella Ulei, 120

Durante, discussion of maize in herbal of, 165

Dwarf mutant in maize, Comparative developmental study of, 289

Ear and tassel in maize: development, 269, 280-286; structure of a dwarf mutant,

Epidermis in Zea Mays, General features of

Euphorbiaceae, living, compared to a fossil wood, 109, 110

Equisetales, relation of, to Sphenophyllum, 218

Extinction of Coleus hairs at various thicknesses, 251

Ferns, fossil, 107

Finan, John J. Maize in the great herbals, 149

Flavone concentration, determination of, in Coleus hair cells, 249, 252

Flora of Panama, Part V, Fasc. 1, 1 Fomes idahoensis Brown, A note on, 207 Food, Uses of Hevea for, in relation to its domestication, 117

Forsteronia, 236; elachista, 237; gracilioides, 237; mediocris, 237; propinqua, 236; spicata, 236

Fossil wood from Idaho, 108

Fossils, fern, 107, 203; in coal balls, 193, 209

Frumentum Asiaticum, 167; Indicum, 160, 167; Turcicum, 159, 167, 173, 177

Fuchs, Leonhard, discussion of maize in herbal of, 159, 172; illustration of maize in, 158, 161

Garuga, 109; pinnata, 110

Gerarde's herbal, description of maize in, 167

Glands, The uses of, in a taxonomic consideration of the family Bignoniaceae, 123 Gómara, description of maize by, 153

Graminese, epidermis in the, 341

cherrima, 143; Ulei, 144

Guilielma Gasipaes, 119 Gynandropsis, 139, 141; adenocarpa, 144; aurantiaca, 143; brachycarpa, 140, 143; chiriquensis, 142; coccinea, 143; decipiens, 142; densiflora, 144; gracilis, 145; birsuta, 144; bispidula, 143; Jamesonii, 144; lateralis, 144; macrophylla, 145; macrothyrsus, 144; Mathewsii, 144; orba, 145; pentaphylla, 139; porphyrantha, 145; phoenicea, 144; puberula, 144; pul-

Herbals, great, Maize in the, 149; passages from original texts quoted and translated, 187; woodcuts of, 179, 184 Hernández, description of maize by, 165

Hernandiaceae of Panama, 68 Hevea, The uses of, for food in relation to

its domestication, 117 Hevea pauciflora, 120

Idaho: coniferous wood found near Wayan, 107; Fomes idahoensis Brown from, 207 Illinois: coal balls from, 204; A Crossotheca from northern, 203

Inflorescence of Zea Mays, 269-406 Iowa, An Osmundaceous stem, from, 193

Jacaranda mimosiflora, capitate glandular hairs on staminodium of, 136

K

Kirkia acuminata, 109, 110 Koordersiodendron, 109; pinnatum, 110

L

Laubengayer, R. A. The vascular anatomy of the four-rowed ear of corn, 337 Lauraceae: living, compared with a fossil

wood, 110; of Panama, 1

Lenz, Lee Wayne. Comparative anatomy of the female inflorescence of Zea Mays L., 353

Light, relation of, to cell investigation, 239 l'Obel, see Obel

Lundia corymbifera, interpetiolar gland on stem of, 134

M

Mahiz, 153

0

ca

3

lar

Maize, see also Corn and Zea Mays

Maize: anatomy of female inflorescence of, 353; Caribbean, 183; cob, 304, 353; colors of kernels according to herbalists, 169; domestication of, 394; dwarf mutant in, 289; ear and tassel development in, 269; epidermis in, 341; evolution of, 377; flints, 182; four-rowed ear, 337; history of, 149, 180, 255; in the great herbals, 149, illustrations, 150-175; in post-conquest Hispanic America, 150; names in herbals for, 168; origin of, 397; place of origin according to herbals, 169; pod corn, role of, in origin of maize, 377; religious ceremonies connected with, 155; row number in ears, 323, 337; southern dent varieties, 255, 268; structure of, 289, 301; tassel and ear development, 269

Malouetia Quadricasarum, 234; tamaquarina, 235; Schomburgkii, 235

Mamay, Sergius, Henry N. Andrews and. A Crossotheca from northern Illinois, 203

Mandevilla longipes, 233; subsagittata, 234; turgida, 234; Vanheurckii, 234

Mangelsdorf, Paul C. The role of pod corn in the origin and evolution of maize, 377 Manisuris cylindrica, 302, 311

Matelea purpureolineata, 237

Matthiolus, description of maize in herbal of, 163, 164; illustration of maize in, 156

Mauria simplicifolia, 110 Memora Klugii, corolla glands of, 132

Mespilodaphne sassafras, 110

Microspectrophotometer, a new: optical diagram of, 240; Quantitative determina-

tion of the pigment content of single cells by means of, 239

Milium indicum, 162

Miscellaneous new Apocynaceae and Asclepiadaceae, 233 Monardes, description of maize in herbal

of, 165

Morangaceae of Panama, 75 Morleya, 233; leipocalyx, 233 Mutant in maize, a dwarf, 289

N

Neuropteris, 204

0

l'Obel, herbal of, 162, 176; illustration of maize in, 163 Origin and evolution of maize, Role of pod

Corn in, 377 Osmundaceae, 193

Osmundaceous stem from Iowa, 193

Osmundites, 195; Dowkeri, 195; Dunlapi, 195; Kolbei, 195; Schemnitzensis, 195; skidegatensis, 196

Oviedo's "Historia natural y general," 150; description of maize in, 151, 154, 164; illustration of maize in, 150

P

Pachyptera Kerere, interpetiolar and petiolar glands of, 128

Paleobotanical studies, 107, 193, 203, 207, 209

Panama, Flora of, Part V, Fasc. 1 (Lauraceae to Cruciferae), 1

Papaveraceae of Panama, 71

Paraphyllanthoxylon arizonense, 108; idahoense, 108, 112, 115

Parkinson, John, discussion of maize in herbal of, 167

Pecopteris, 203, 204

Pedicellaris, 139; Lebmannii, 144; Ulei, 144 Petitia domingensis, 110, 111

Phyllanthus emblica, 109, 110

Pigment content of single cells, Quantitative determination of, by means of a new microspectrophotometer, 239

Plumeria, 233

Pod corn, Role of, in the origin and evolution of maize, 377

Podandrogyne, 139, 141; brachycarpa, 143; cernua, 143; chiriquensis, 142, 146; coccinea, 143, 146; decipiens, 142; glabra, 141, 145; gracilis, 144; macrophylla, 145; orba, 145; polychroma, 145; pubescens, 145

Polypore, A fossil, from Idaho, 207 Prat. Henri. General features of the epidermis in Zea Mays, 341

Prestonia didyma, 236; Haughtii, 235; macrophylla, 235; obovata, 236 Protoosmundites Wilsonii, 194, 198, 201 Pseudocalymma sp., interpetiolar glands of,

Pteridosperm stems in Iowa coal balls, 193

0

Quantitative determination of the pigment content of single cells by means of a new microspectrophotometer, 239

Religious ceremonies, maize in, 155

Rhodea, 204

Right-handed and left-handed corn embryos, 317

Rollins, Reed C. Cruciferae of Panama, 99 Row number in maize, A morphological analysis of, 323

Schery, Robert W., Robert E. Woodson, Jr., and, and collaborators. Flora of Panama. Part V. Fascicle 1 (Lauraceae to Cruciferae), 1

Schinopsis balansea, 110

Seibert, R. J.: The use of glands in a taxonomic consideration of the family Bignoniaceae, 123; The uses of Hevea for food in relation to its domestication, 117 Simarubaceae, living, compared to a fossil

wood, 109, 110

Siphonia, 118

Southern dent corns, 255, 268; chromosome knobs, 257, 258; internode diagram, 260, 261; old southern dents, 262; derived varieties, 263

Spackman, William, Jr. A dicotyledonous wood found associated with the Idaho

Tempskyas, 107

Spectrophotometric measurements, determination of anthocyanin and flavone concentration in cells by, 239, 242

Sphenophyllum, Vegetative anatomy of, 209

Sphenophyllum, 209; Dawsoni, 209; Gilkineti, 210, 218; insigne, 210, 217; minus, 218; perforatum, 210, 217; plurifoliarum, 210, 217, 220-231

Sphenopteris 203

Staphylopteris sagittatus, 204

Stem remains of ferns in coal balls, 193 Stephens, S. G. A comparative developmental study of a dwarf mutant in maize and its bearing on the interpretation of tassel and ear structure, 289

Tassel in maize: development of, 269, 280-286; structure of, in a dwarf mutant, 289 Tempskyas, Idaho, A dicotyledonous wood

found associated with, 107

Tabernaemontanus' herbal, description of maize in, 160, 177; illustrations of maize in, 173-177

Teopod maize ear, 310, 310

Teosinte, 302, 311, 317

Thamnopteris, 196; Schlechtendalii, 196 Tlaolli, 165

Tovariaceae of Panama, 75

Tripsacum, 302, 311

Triticum Bactrianum, 160; Turcicum, 162

U

Urticaceae, 111

Vascular anatomy of the four-rowed ear of corn, 337

Verbenaceae, living, compared to a fossil wood, 110

Weatherwax, Paul. Right-handed and lefthanded corn embryos, 317

Welschen Korn, 156

Woodcuts: botanical, 185; copying, 186; historical background of, 184; of maize

in the herbals, 170, 179

Woodson, Robert E., Jr.: Gynandropsis, Cleome, and Podandrogyne, 139; Miscellaneous new Apocynaceae and Asclepiadaceae, 233; and Robert W. Schery and collaborators. Flora of Panama. Part V. Fascicle 1 (Lauraceae to Cruciferae), 1

Zalesskya, 195; diploxylon, 196; gracilis, 196

Zea Mays-See Corn and Maize

Zea Mays L., Comparative histology of the female inflorescence of, 353; General features of the epidermis in, 341

Zea mexicana, 302

